

H2 MATHS

Exam papers with worked solutions

SET B

PAPER 1

ANSWER

Compiled by

THE MATHS CAFE

Qns	Solution	Notes
1	<p>Let the amount of money for a blue voucher, a yellow voucher and a red voucher be x, y and z respectively.</p> <p>Then</p> $3x + 4y + 7z = 27.40 \quad \dots\dots(1)$ $5x + 2y + 4z = 20.80 \quad \dots\dots(2)$ $2x + 8y + 5z = 45.00 \quad \dots\dots(3)$ <p>From G.C., $x = 2$, $y = 5$, $z = 0.20$</p> $4x + py + 2z = 43.40$ $p = \frac{43.40 - 4(2) - 2(0.2)}{5} = 7$	
2 (i)	<p>Let P_n be the statement $u_n = \frac{1}{3}[1 + 8(-2)^n]$ for all $n \in \mathbb{N}_0^+$.</p> <p>LHS of $P_0 = u_0 = 3$ (Given)</p> <p>RHS of $P_0 = \frac{1}{3}[1 + 8(-2)^0] = 3$</p> <p>$\therefore P_0$ is true.</p> <p>Assume that P_k is true for some $k \in \mathbb{N}_0^+$, ie</p> $u_k = \frac{1}{3}[1 + 8(-2)^k].$ <p>We want to prove P_{k+1} is true, ie $u_{k+1} = \frac{1}{3}[1 + 8(-2)^{k+1}]$.</p> <p>LHS of $P_{k+1} = u_{k+1} = 1 - 2u_k$ (Given)</p> $= 1 - \frac{2}{3}[1 + 8(-2)^k]$ $= \frac{1}{3} - \frac{16}{3}(-2)^k$ $= \frac{1}{3} + \frac{8}{3}(-2)(-2)^k$ $= \frac{1}{3}[1 + 8(-2)^{k+1}]$ <p>$\therefore P_{k+1}$ is true</p> <p>Since P_0 is true and P_k is true $\Rightarrow P_{k+1}$ is true, by Mathematical Induction, P_n is true for all $n \in \mathbb{N}_0^+$.</p>	
(ii)	<p>The sequence is divergent as $n \rightarrow \infty$, $(-2)^n$ does not converge to a finite number</p>	

<p>3</p>	<p>when $r = 0$, $3 = 3b$ $r = -1$, $1 = (4 - a)b$</p>	
	<p>Alternative Method $(r + 1)^4 + (r + 1)^2 + 1 \equiv (r^2 + ar + 3)(r^2 + r + b)$ $(r^4 + 4r^3 + 6r^2 + 4r + 1) + (r^2 + 2r + 1) + 1 \equiv$ $(r^2 + ar + 3)(r^2 + r + b)$ Comparing coefficients: r^0: $3 = 3b$ r^3: $4 = 1 + a$</p>	
	<p>$a = 3, b = 1$</p>	
	$\frac{1}{r^2 + r + 1} - \frac{1}{r^2 + 3r + 3} = \frac{(r^2 + 3r + 3) - (r^2 + r + 1)}{(r^2 + r + 1)(r^2 + 3r + 3)}$ $= \frac{2(r + 1)}{(r + 1)^4 + (r + 1)^2 + 1}$	
	$\sum_{r=0}^N \frac{r + 1}{(r + 1)^4 + (r + 1)^2 + 1}$ $= \frac{1}{2} \sum_{r=0}^N \left(\frac{1}{r^2 + r + 1} - \frac{1}{r^2 + 3r + 3} \right)$ $= \frac{1}{2} \left[\begin{array}{c} \frac{1}{1} - \frac{1}{3} \\ \cancel{\frac{1}{3}} - \cancel{\frac{1}{7}} \\ \vdots \\ \frac{1}{(N-1)^2 + (N-1) + 1} - \frac{1}{(N-1)^2 + 3(N-1) + 3} \\ \cancel{\frac{1}{N^2 + N + 1}} - \cancel{\frac{1}{N^2 + 3N + 3}} \end{array} \right]$ $= \frac{1}{2} \left(1 - \frac{1}{N^2 + 3N + 3} \right)$	
	$\sum_{r=2}^N \frac{r}{r^4 + r^2 + 1} = \sum_{r=1}^{N-1} \frac{r + 1}{(r + 1)^4 + (r + 1)^2 + 1}$ $= \frac{1}{2} \left(\frac{1}{3} - \frac{1}{(N-1)^2 + 3(N-1) + 3} \right)$ $= \frac{1}{2} \left(\frac{1}{3} - \frac{1}{N^2 + N + 1} \right)$	

<p>4 (i)</p>	<p>Area of region R</p> $= 5 \times \sqrt{3} - \int_0^{\sqrt{3}} \left(\frac{2}{\sqrt{4-x^2}} + 3 \right) dx$ $= 1.37 \quad (\text{to 3 s.f.})$	
<p>(ii)</p>	<p>Equation of new curve</p> $y = \frac{2}{\sqrt{4-x^2}} + 3 - 5$ $y = \frac{2}{\sqrt{4-x^2}} - 2$	
	<p>Volume of revolution</p> $= \pi \int_0^{\sqrt{3}} \left(\frac{2}{\sqrt{4-x^2}} - 2 \right)^2 dx$ $= \pi \int_0^{\sqrt{3}} \left(\frac{4}{4-x^2} - \frac{8}{\sqrt{4-x^2}} + 4 \right) dx$ $= 4\pi \int_0^{\sqrt{3}} \left(\frac{1}{4-x^2} - \frac{2}{\sqrt{4-x^2}} + 1 \right) dx \quad (\text{Shown})$ $= 4\pi \left[\frac{1}{2(2)} \ln \left \frac{2+x}{2-x} \right - 2 \sin^{-1} \frac{x}{2} + x \right]_0^{\sqrt{3}}$ $= 4\pi \left[\frac{1}{4} \ln \left \frac{2+\sqrt{3}}{2-\sqrt{3}} \right - \frac{2\pi}{3} + \sqrt{3} \right]$	
<p>5</p>	$\pi r l = k\pi$ $l = \frac{k}{r}$ $l^2 = r^2 + h^2$ $h = \sqrt{\frac{k^2}{r^2} - r^2} = \frac{\sqrt{k^2 - r^4}}{r}$ $V = \frac{1}{3} \pi r^2 \left(\frac{\sqrt{k^2 - r^4}}{r} \right)$ $V = \frac{\pi r \sqrt{k^2 - r^4}}{3}$ <hr/> $\frac{dV}{dr} = \frac{\pi \sqrt{k^2 - r^4}}{3} - \frac{2\pi r^4}{3\sqrt{k^2 - r^4}}$	

	<p>At stationary point, $\frac{dV}{dr} = 0$</p> $\frac{\pi\sqrt{k^2 - r^4}}{3} = \frac{2\pi r^4}{3\sqrt{k^2 - r^4}}$ $k^2 - r^4 = 2r^4$ $3r^4 = k^2$ $r = \sqrt[4]{\frac{k^2}{3}}$	
--	---	--

6	$u = \ln x \quad \frac{dv}{dx} = \frac{1}{x^2}$	
(a)	$\frac{du}{dx} = \frac{1}{x} \quad v = -\frac{1}{x}$	
(i)		

	$\int_1^n \frac{1}{x^2} \ln x \, dx$ $= \left[-\frac{1}{x} \ln x \right]_1^n - \int_1^n -\frac{1}{x} \left(\frac{1}{x} \right) dx$ $= -\frac{\ln n}{n} - \left[\frac{1}{x} \right]_1^n$ $= -\frac{\ln n}{n} - \left[\frac{1}{n} - 1 \right]$ $= -\frac{\ln n}{n} - \frac{1}{n} + 1$	
--	---	--

(a)	$\int_1^\infty \frac{1}{x^2} \ln x \, dx = \lim_{n \rightarrow \infty} \left[-\frac{\ln n}{n} - \frac{1}{n} + 1 \right]$	
(ii)	$= 1$	


(b)	$x = a \sec \theta \Rightarrow \frac{dx}{d\theta} = a \sec \theta \tan \theta$ <p>When $x = a$, $\sec \theta = 1 \Rightarrow \cos \theta = 1 \Rightarrow \theta = 0$.</p> <p>When $x = 2a$, $\sec \theta = 2 \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$.</p>	
------------	---	--

	$\int_a^{2a} \frac{\sqrt{x^2 - a^2}}{x} dx$ $= \int_0^{\frac{\pi}{3}} \frac{\sqrt{a^2 \sec^2 \theta - a^2}}{a \sec \theta} a \sec \theta \tan \theta d\theta$ $= a \int_0^{\frac{\pi}{3}} \tan^2 \theta d\theta$ $= a \int_0^{\frac{\pi}{3}} (\sec^2 \theta - 1) d\theta$ $= a [\tan \theta - \theta]_0^{\frac{\pi}{3}}$ $= a \left(\sqrt{3} - \frac{\pi}{3} \right)$	
--	--	--

7	Let A_n denote the distance ran on the n th training session and S_n denote the total distance ran for the n training sessions.	
(i)	$A_n = 7.5 + 0.8(n - 1)$ $= 6.7 + 0.8n$	
(ii)	$S_n \geq 475$ $\frac{n}{2} [2(7.5) + 0.8(n - 1)] \geq 475$ $7.1n + 0.4n^2 \geq 475$ From GC, $n = 26.7$ Least $n = 27$	
(iii)	For the modified training session, let B_n denote the distance ran on the n th training session and G_n denote the total distance ran for the n training sessions.	
	$B_6 = 14.93$ $x(1.2)^5 = 14.93$ $x = 6 \text{ (nearest integer)}$	

(iv)	$G_n = \frac{6\left[\left(\frac{6}{5}\right)^n - 1\right]}{\frac{6}{5} - 1}$ $= 30\left[\left(\frac{6}{5}\right)^n - 1\right]$	
	$\sum_{n=1}^N G_n = \sum_{n=1}^N 30\left[\left(\frac{6}{5}\right)^n - 1\right]$ $= 30\sum_{n=1}^N \left(\frac{6}{5}\right)^n - \sum_{n=1}^N 30$ $= \frac{30 \cdot \frac{6}{5} \left[\left(\frac{6}{5}\right)^N - 1\right]}{\frac{6}{5} - 1} - 30N$ $= 6\left\{30\left[\left(\frac{6}{5}\right)^N - 1\right]\right\} - 30N$ $= 6G_N - 30N$	


8 (i)	$y = \frac{-x^2 + 4x - 5}{x - 2}$ $\Rightarrow y(x - 2) = -x^2 + 4x - 5$ $\Rightarrow x^2 + (y - 4)x + (5 - 2y) = 0$ <p>For real values of x, discriminant ≥ 0.</p> $\Rightarrow (y - 4)^2 - 4(5 - 2y) \geq 0$ $\Rightarrow y^2 \geq 4$ $\Rightarrow y \leq -2 \text{ or } y \geq 2$	
(ii)	$\frac{-x^2 + 4x - 5}{x - 2} = A(x - 2) + \frac{B}{x - 2}$ $\Rightarrow -x^2 + 4x - 5 = A(x - 2)^2 + B$ <p>Comparing coefficients of x^2, $A = -1$ Comparing constants, $-5 = 4A + B \Rightarrow B = -1$</p>	
	$y = \frac{-x^2 + 4x - 5}{x - 2} = -(x - 2) - \frac{1}{x - 2}$ $y = x + \frac{1}{x} \xrightarrow{T_1} y = x - 2 + \frac{1}{x - 2} \xrightarrow{T_2} y = -\left(x - 2 + \frac{1}{x - 2}\right)$	
	<p>The graph of $y = x + \frac{1}{x}$ can be transformed to the graph of $y = \frac{-x^2 + 4x - 5}{x - 2}$ using the following transformations in succession.</p> <p>T_1: Translation of 2 units in the positive x direction. T_2: Reflection in the x-axis.</p>	




9(i)	least $a = 4$	
(ii)	<p>when $x > 4$, $f(x) = y = \frac{2x-8}{x-2}$</p> $xy - 2y = 2x - 8$ $x(y - 2) = 2y - 8$ $x = \frac{2y - 8}{y - 2}$	
	$f^{-1} : x \mapsto \frac{2x-8}{x-2}, 0 < x < 2$	
(iii)	<p>$g(x) = x^2 - 6x + 7 = (x - 3)^2 - 2, x < 3$</p> <p>$D_g = (-\infty, 3)$ $D_f = (4, \infty)$ $R_g = (-2, \infty)$ $R_f = (0, 2)$</p>	
	<p>fg does not exist since $R_g = (-2, \infty) \not\subseteq (4, \infty) = D_f$</p> <p>$gf$ exists since $R_f = (0, 2) \subseteq (-\infty, 3) = D_g$</p>	

	$gf(x) = g\left(\frac{2x-8}{x-2}\right)$ $= \left(\frac{2x-8}{x-2}\right)^2 - 6\left(\frac{2x-8}{x-2}\right) + 7, x > 4$	
	From GC: $R_{gf} = (-1, 7)$	
	Alternative method (Mapping method): $(4, \infty) \xrightarrow{f} (0, 2) \xrightarrow{g} (-1, 7)$ $R_{gf} = (-1, 7)$	

10	Given A, B and C are collinear,	
(i)	$\vec{AC} = k\vec{AB}$ $\mathbf{c} - \mathbf{a} = k(\mathbf{b} - \mathbf{a})$ $\mathbf{c} = k\mathbf{b} + (1 - k)\mathbf{a}$ (shown)	
(ii)	$ \mathbf{a} \times \mathbf{c} = \mathbf{a} \times [k\mathbf{b} + (1 - k)\mathbf{a}] $ $= k(\mathbf{a} \times \mathbf{b}) + (1 - k)(\mathbf{a} \times \mathbf{a}) $ $= k \mathbf{a} \mathbf{b} \sin 90^\circ \hat{\mathbf{n}} + (1 - k)\mathbf{0} $ $= 9 k $	
	It is the area of a parallelogram with sides OA and OC .	
(iii)	Area of triangle $OAC = 3 \times$ area of triangle OAB $\frac{1}{2} \mathbf{a} \times \mathbf{c} = \frac{3}{2} \mathbf{a} \times \mathbf{b} $ $9 k = 3 \mathbf{a} \mathbf{b} \sin 90^\circ$ $= 27$ $ k = 3$ $k = \pm 3$	
(iv)	Length of projection of OC onto $OA = 12$ $\frac{ \mathbf{c} \cdot \mathbf{a} }{ \mathbf{a} } = 12$ $ \mathbf{c} \cdot \mathbf{a} = 12 \mathbf{a} $ $= 36$	
	When $k = 3, \mathbf{c} = 3\mathbf{b} - 2\mathbf{a}$ $ \mathbf{c} \cdot \mathbf{a} = (3\mathbf{b} - 2\mathbf{a}) \cdot \mathbf{a} $ $= 3(\mathbf{b} \cdot \mathbf{a}) - 2\mathbf{a} \cdot \mathbf{a} $ $= 3(0) - 2(3)^2 $ $= 18$	
	When $k = -3, \mathbf{c} = -3\mathbf{b} + 4\mathbf{a}$ $ \mathbf{c} \cdot \mathbf{a} = (-3\mathbf{b} + 4\mathbf{a}) \cdot \mathbf{a} $ $= -3(\mathbf{b} \cdot \mathbf{a}) + 4\mathbf{a} \cdot \mathbf{a} $ $= -3(0) + 4(3)^2 $ $= 36$	
	$\therefore \mathbf{c} = -3\mathbf{b} + 4\mathbf{a}$	

<p>11 (a)</p>	$z = (1+i)t + \frac{1-i}{t}$ $= \left(t + \frac{1}{t}\right) + \left(t - \frac{1}{t}\right)i$ <p>Let $x = t + \frac{1}{t}$ -----(1)</p> $y = t - \frac{1}{t}$ -----(2) <p>(1) + (2): $x + y = 2t$</p> <p>(1) - (2): $x - y = \frac{2}{t}$</p> $\therefore x - y = \frac{2}{\frac{x+y}{2}}$ $\Rightarrow x^2 - y^2 = 4$	
<p>11 (b)</p>		
	<p>If $\arg(p) > \arg(q)$,</p> $\arg\left(\frac{p}{q}\right) = \arg p - \arg q = \angle POQ$ $= 2 \tan^{-1} \frac{2 a }{3 a }$ $= 2 \tan^{-1} \frac{2}{3} \text{ or } 1.18 \text{ rad}$	
	$ p+q = 2 3a = 6 a $	

<p>12 (i)</p>	$x = t^2, \quad y = t^3 - 4$ $\frac{dx}{dt} = 2t, \quad \frac{dy}{dt} = 3t^2$ $\frac{dy}{dx} = \frac{3t}{2}$	
	<p>Tangent at P</p> $(y - p^3 + 4) = \frac{3p}{2}(x - p^2)$ $2y = 3px - p^3 - 8$	
<p>(ii)</p>	<p>Since the tangent passes through the origin, subst. $x = 0$ and $y = 0$ into the equation of tangent in part (i).</p> $-p^3 - 8 = 0$ $p = -2$ $x = 4, y = -12$ $P(4, -12)$	
<p>(iii)</p>	<p>$x = 0, y = -4, t = 0$</p> 	
<p>(iv)</p>	$\text{Area} = \frac{1}{2}(4)(12) - \int_{-12}^{-4} x \, dy$ $= 24 - \int_{-2}^0 (t^2)(3t^2) \, dt$ $= 24 - \left[\frac{3t^5}{5} \right]_{-2}^0$ $= \frac{24}{5}$	