

# **H2 MATHS**

**Exam papers with worked solutions**


## **SET B**

## **PAPER 2**


## **ANSWER**

Compiled by

**THE MATHS CAFE**

Qn	Solution	Notes
1(a) (i)	$e^{y+x} = \cos x$ $y = \ln(\cos x) - x$ Differentiate wrt $x$ $\frac{dy}{dx} = -\tan x - 1$ Differentiate wrt $x$ $\frac{d^2y}{dx^2} = -\sec^2 x$ $x = 0$ $y = 0$ $\frac{dy}{dx} = -1$ $\frac{d^2y}{dx^2} = -1$ $y = -x - \frac{x^2}{2} + \dots$	
(a) (ii)	$ h(x) - y  < 0.2$ $ h(x) - y  - 0.2 < 0$ <div style="border: 1px solid black; padding: 5px; margin: 5px 0;"> <pre> Plot1 Plot2 Plot3 \Y1=-X-X^2/2 \Y2=ln(cos(X))-X \Y3= Y1-Y2 -0.2 \Y4= \Y5= Zero X=-1.116117 Y=0 Zero X=1.1161165 Y=0                     </pre>  </div> From GC, $-1.12 < x < 1.12 \text{ (to 3 s.f.)}$	

1(b)	$\left(a - \frac{x}{3}\right)^n = a^n \left(1 - \frac{x}{3a}\right)^n$	
	$= a^n \left(1 - \frac{xn}{3a} + \frac{n(n-1)}{2} \left(\frac{x}{3a}\right)^2 + \dots\right)$ $= a^n \left(1 - \frac{n}{3a}x + \frac{n(n-1)}{18a^2}x^2 + \dots\right)$	
	$-\frac{n}{3a} = \frac{4n(n-1)}{18a^2}$ $n = 1 - \frac{3a}{2}$	
	$a^n = \frac{1}{4}$ <p>Sub <math>n = 1 - \frac{3a}{2}</math></p> $a^{1 - \frac{3a}{2}} = \frac{1}{4}$	
	<p>From GC, <math>a = 2</math> or <math>0.16086</math> (rejected)</p> <p>when <math>a = 2</math>, <math>n = -2</math></p>	

<p><b>2(a)</b></p>	$\frac{d^2y}{dx^2} = ae^{-2x}$ $\frac{dy}{dx} = \frac{ae^{-2x}}{-2} + c$ $y = \frac{ae^{-2x}}{4} + cx + d$	
<p><b>2(b)</b></p>	$\frac{dx}{dt} = kx - p$ <p>Given that <math>\frac{dx}{dt} = 0</math> when <math>x = 12</math>.</p> $\Rightarrow 12k - p = 0$ $\Rightarrow k = \frac{p}{12}$ $\therefore \frac{dx}{dt} = \frac{px}{12} - p$ $\Rightarrow \frac{dx}{dt} = \frac{p}{12}(x-12) \quad (\text{Shown})$	
<p><b>(i)</b></p>	$\int \frac{1}{x-12} dx = \int \frac{p}{12} dt$ $\ln x-12  = \frac{p}{12}t + c$ $x-12 = \pm e^{\frac{p}{12}t + c}$ $x = 12 + Ae^{\frac{p}{12}t}, \quad \text{where } A = \pm e^c$ <p>When <math>t = 0</math>, <math>x = 10 \Rightarrow A = -2</math></p> $\therefore x = 12 - 2e^{\frac{p}{12}t}$	
<p><b>(ii)</b></p>	<p>When <math>x = 0</math>, <math>t = T</math></p> $\Rightarrow 12 - 2e^{\frac{p}{12}T} = 0$ $\Rightarrow \frac{p}{12}T = \ln 6$ $\Rightarrow T = \frac{12}{p} \ln 6$	
<p><b>(c)</b></p>		

3	(i)	<p>Since <math>A</math> lies in the plane <math>\pi_1</math>, <math>\begin{pmatrix} -10 \\ 0 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} \alpha \\ \beta \\ 0 \end{pmatrix} = -30</math>.</p> <p><math>\Rightarrow -10\alpha = -30</math>  <math>\Rightarrow \alpha = 3</math></p>	
		<p>Perpendicular distance from <math>O</math> to <math>\pi_1 = 6</math></p> <p><math>\Rightarrow \frac{ \overrightarrow{OA} \cdot \mathbf{n}_1 }{ \mathbf{n}_1 } = 6</math></p> <p><math>\Rightarrow \frac{ -30 }{ \mathbf{n}_1 } = 6</math></p> <p><math>\Rightarrow  \mathbf{n}_1  = 5</math>  <math>\Rightarrow \alpha^2 + \beta^2 = 5^2</math>  <math>\Rightarrow \beta = 4</math></p>	
	(ii)	<p>Acute angle between <math>OA</math> and <math>\pi_2</math></p> <p><math>= \sin^{-1} \frac{ \overrightarrow{OA} \cdot \mathbf{n}_2 }{ \overrightarrow{OA}   \mathbf{n}_2 }</math></p> <p><math>= \sin^{-1} \frac{\left  \begin{pmatrix} -10 \\ 0 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} \right }{\left  \begin{pmatrix} -10 \\ 0 \\ 5 \end{pmatrix} \right  \left  \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} \right }</math></p> <p><math>= \sin^{-1} \frac{20}{\sqrt{125} \sqrt{6}}</math>  <math>= 46.9^\circ</math></p>	
	(iii)	<p><math>\begin{pmatrix} -10 \\ 0 \\ 5 \end{pmatrix} \times \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -5 \\ 15 \\ -10 \end{pmatrix}</math></p> <p>A normal to plane <math>\pi_3</math> is <math>\begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix}</math>.</p> <p>A cartesian equation is <math>x - 3y + 2z = 0</math>.</p>	


<p><b>4(i)</b></p>	$p = 1 - i, \quad q = 1 + \sqrt{3}i, \quad r = k + i$ $\Rightarrow  p  = \sqrt{1^2 + (-1)^2} = \sqrt{2}, \quad  q  = \sqrt{1^2 + (\sqrt{3})^2} = 2, \quad  r  = \sqrt{k^2 + 1}$ $\arg(p) = -\frac{1}{4}\pi, \quad \arg(q) = \frac{1}{3}\pi$ $ s  = \frac{1}{2}$ $\Rightarrow \frac{ p ^2  r }{ q ^3} = \frac{1}{2}$ $\Rightarrow \frac{(\sqrt{2})^2 \sqrt{k^2 + 1}}{(2)^3} = \frac{1}{2}$	
	$\Rightarrow \sqrt{k^2 + 1} = 2$ $\Rightarrow k^2 + 1 = 4$ $\Rightarrow k = \pm\sqrt{3}$	
	$\arg(s) = -\frac{2}{3}\pi$ $\Rightarrow 2\arg(p) + \arg(r) - 3\arg(q) = -\frac{2}{3}\pi$ $\Rightarrow \arg(r) = -\frac{2}{3}\pi - 2\left(-\frac{1}{4}\pi\right) + 3\left(\frac{1}{3}\pi\right) = \frac{5}{6}\pi$	
	<p>If <math>k = \sqrt{3}</math>, then <math>\arg(r) = \frac{1}{6}\pi</math></p> <p>If <math>k = -\sqrt{3}</math>, then <math>\arg(r) = \frac{5}{6}\pi</math></p> <p><math>\therefore k = -\sqrt{3}</math></p>	

<p><b>4(ii)</b></p>	$z^3 = 4(1 + \sqrt{3}i)$ $z^3 = 8e^{i\left(\frac{1}{3}\pi + 2k\pi\right)}$ $= 8e^{i\left(\frac{1+6k}{3}\right)\pi}$	
	$z = 2e^{i\left(\frac{1+6k}{9}\right)\pi}, \quad k = -1, 0, 1$ $= 2e^{i\left(-\frac{5}{9}\pi\right)}, \quad 2e^{i\left(\frac{\pi}{9}\right)}, \quad 2e^{i\left(\frac{7}{9}\pi\right)}$	
	$w^3 = 4(\sqrt{3} - i)$ $w^3 = 4i(-\sqrt{3}i - 1)$ $w^3 = -4i(1 + \sqrt{3}i)$ $w^3 = (i)^3 4(1 + \sqrt{3}i) \text{ since } i^2 = -1$ $\left(\frac{w}{i}\right)^3 = 4(1 + \sqrt{3}i)$	
	<p>Comparing with <math>z^3 = 4(1 + \sqrt{3}i)</math></p> $\frac{w}{i} = z$ $w = iz$	
	$w = e^{i\left(\frac{\pi}{2}\right)} z$ $= e^{i\left(\frac{\pi}{2}\right)} \times 2e^{i\left(-\frac{5}{9}\pi\right)}, \quad e^{i\left(\frac{\pi}{2}\right)} \times 2e^{i\left(\frac{\pi}{9}\right)}, \quad e^{i\left(\frac{\pi}{2}\right)} \times 2e^{i\left(\frac{7}{9}\pi\right)}$ $= 2e^{i\left(-\frac{1}{18}\right)\pi}, \quad 2e^{i\left(\frac{11}{18}\right)\pi}, \quad 2e^{i\left(\frac{13}{18}\right)\pi}$	


<b>5(i)</b>	This is quota sampling.		
	A disadvantage of quota sampling is that the sample is not a good representation of the population (residents of the town).		
<b>(ii)</b>	Select the following number of students <u>randomly</u> from each stratum below.		
		<b>18 - 25 year old</b>	<b>26 - 35 year old</b>
	<b>Male</b>	$\frac{75}{250} \times 100 \approx 30$	$\frac{36}{250} \times 100 \approx 14$
	<b>Female</b>	$\frac{99}{250} \times 100 \approx 40$	$\frac{40}{250} \times 100 \approx 16$

<b>6</b>	Given $X \sim N(\mu, \sigma^2)$ , $P(2X_1 < X_2) = 0.8$ $P(2X_1 - X_2 < 0) = 0.8$ Now, $2X_1 - X_2 \sim N(\mu, 5\sigma^2)$
	$P\left(Z < -\frac{\mu}{\sqrt{5}\sigma}\right) = 0.8$ $-\frac{\mu}{\sqrt{5}\sigma} = 0.84162$
	$\mu = -1.8819\sigma$ $\sigma = -0.53138\mu$
	$P(X_1 + X_2 > 2a) = 0.8$ $P(X_1 + X_2 < 2a) = 0.2$ Now, $X_1 + X_2 \sim N(2\mu, 2\sigma^2)$
	$P\left(Z < \frac{2a - 2\mu}{\sqrt{2}\sigma}\right) = 0.2$ $\frac{2a - 2\mu}{\sqrt{2}\sigma} = -0.84162$
	$a - \mu = -0.59512\sigma$ $a - \mu = -0.59512(-0.53138\mu)$ $\mu = 0.75974a$ $= 0.760a$



7(i)	$P(B A') = 0.75$ $\frac{P(B \cap A')}{P(A')} = 0.75$	
	$P(B \cap A') = 0.75(1 - 0.6)$ $= 0.3$ 	
	$P(A' \cap B') = 1 - P(A) - P(B \cap A')$ $= 1 - 0.6 - 0.3$ $= 0.1$	
(ii)	$P(A B) = 0.4$ $\frac{P(A \cap B)}{P(B)} = 0.4$ $P(A \cap B) = 0.4P(B)$ $P(B) = P(B \cap A') + P(A \cap B)$ $= 0.3 + 0.4P(B)$ $\therefore 0.6P(B) = 0.3$ $P(B) = 0.5$	
	$P(A) = 0.6$ $P(A B) = 0.4$ $\therefore P(A B) \neq P(A)$ $\therefore A \text{ and } B \text{ are not independent.}$	

<b>8</b>	1A, 1B, 1S, 2E, 1N, 1C No. of code words formed with 2 'E's = ${}^5C_2 \times \frac{4!}{2!}$ = 120	
	No. of code words formed with 1 or no 'E' = ${}^6C_4 \times 4!$ = 360	
	Total no. of code words = $120 + 360$ = 480 (shown)	
<b>(i)</b>	$P(\text{four-letter code words contain distinct letters})$ $= \frac{\text{No. of code words formed with 1 or no 'E'}}{\text{No. of code words formed without restrictions}}$ $= \frac{360}{480}$ $= 0.75$	
<b>(ii)</b>	$P\left(\begin{array}{l} \text{code words do not contain} \\ \text{any vowels} \end{array} \middle  \begin{array}{l} \text{code words contain} \\ \text{distinct letters} \end{array}\right)$ $= \frac{\binom{{}^4C_4 \times 4!}{480}}{0.75}$ $= \frac{1}{15}$	

<p><b>9(i)</b></p>	$y = -0.175x + 3.57$ $\bar{y} = -0.175\bar{x} + 3.57$ <hr/> $\frac{22.94 + k}{9} = -0.175\left(\frac{38.8}{9}\right) + 3.57$ <hr/> $\therefore k = 2.40 \text{ (shown)}$	
<p><b>(ii)</b></p>	 <hr/> <p>Using GC: <math>r = -0.943</math> (3 s.f.)</p> <p>Even though <math>r</math> is close to <math>-1</math>, from the scatter diagram, the data points do not follow a straight line. Therefore, a linear model may not be suitable.</p>	
<p><b>(iii)</b></p>	<p>C is the appropriate model as the data points in the scatter diagram follow the graph of <math>y = e + \frac{f}{x}</math>.</p>	
<p><b>(iv)</b></p>	<p>Equation of regression line</p> $y = \frac{2.7480}{x} + 2.0651$ $y = \frac{2.75}{x} + 2.07 \text{ (to 3 s.f.)}$ <hr/> <p>When <math>x = 8.0</math>,</p> $y = \frac{2.7480}{8.0} + 2.0651 = 2.41 \text{ (to 2 d.p.)}$ <hr/> <p>Since <math>x = 8.0</math> falls outside the data range of <math>x</math>, the estimation of <math>y</math> is unreliable.</p>	

<p><b>10(i)</b></p>	<p><math>\sum x = 1620</math> and <math>\sum x^2 = 221175</math></p> <p>Unbiased estimate of population mean = <math>\frac{1620}{12}</math> = 135</p> <p>Unbiased estimate of population variance = <math>\frac{1}{11} \left[ 221175 - \frac{1620^2}{12} \right]</math> = 225</p>	
<p><b>(ii)</b></p>	<p><math>H_0: \mu = 124.5</math> <math>H_1: \mu \neq 124.5</math></p> <p>Let <math>X</math> = number of packets of cereal sold daily. Since <math>n = 12</math> is small,</p> <p>Under <math>H_0</math>, <math>T = \frac{\bar{X} - \mu_0}{\sqrt{\frac{S^2}{n}}} \sim t(n-1)</math></p> <p><math>\alpha = 0.01</math> From GC, <math>p</math>-value = 0.0337</p> <p>Since <math>p</math>-value = 0.0337 &gt; <math>\alpha = 0.01</math>, we do not reject <math>H_0</math> at 1% level of significance and conclude there is insufficient evidence that the mean number of packets of cereal sold per day has changed.</p>	
<p><b>(iii)</b></p>	<p><math>H_0: \mu = 124.5</math> <math>H_1: \mu &gt; 124.5</math></p> <p>Under <math>H_0</math>, <math>\bar{X} \sim N\left(124.5, \frac{13^2}{12}\right)</math></p> <p>Test statistic: <math>z = \frac{\bar{x} - 124.5}{\sqrt{\frac{169}{12}}}</math></p> <p><math>\alpha = 0.01</math> Reject <math>H_0</math> if <math>z \geq 2.3263</math></p> <p>Since <math>H_0</math> is rejected, <math>\frac{\bar{x} - 124.5}{\sqrt{\frac{13^2}{12}}} \geq 2.3263</math> <math>\bar{x} \geq 133.23</math> <math>\bar{x} \geq 133</math></p> <p>It means that there is a probability of 0.01 of concluding that the mean number of packets of cereal sold per day is more than 124.5, when in fact the mean number is 124.5.</p>	

<p><b>11</b> <b>(i)</b></p>	<p>Let <math>X</math> be the number of emergency admissions per day  <math>X \sim \text{Po}(3)</math>  <math>X_1 + X_2 \sim \text{Po}(6)</math>  <math>P(X_1 + X_2 = 3) = 0.0892</math> ( to 3 s.f. )</p>	
<p><b>(ii)</b></p>	<p>Let length of time = <math>t</math> hr  Let <math>T</math> be the number of emergency admissions in <math>t</math> hours.  <math>T \sim \text{Po}\left(\frac{3t}{24}\right)</math>  <math>P(T = 0) = e^{-\frac{3t}{24}} = 0.2</math>  <math>-\frac{3t}{24} = \ln 0.2</math>  <math>t = -8 \ln 0.2</math>  <math>\approx 13 \text{ hr}</math></p>	
<p><b>(iii)</b></p>	<p><math>P(X &gt; 4) = 1 - P(X \leq 4)</math>  <math>= 0.18474 \approx 0.185</math> ( to 3 s.f. )</p>	
<p><b>(iv)</b></p>	<p>Let <math>Y</math> be the number of days with at most 4 admissions out of 50 days.  <math>Y \sim \text{B}(50, 0.81526)</math>  Since <math>n = 50</math> is large, <math>np = 40.763 &gt; 5</math>, <math>nq = 9.237 &gt; 5</math>,  <math>Y \sim \text{N}(40.763, 7.5306)</math> approximately  <math>P(Y &lt; 40) \xrightarrow{\text{c.c.}} P(Y &lt; 39.5) \approx 0.323</math></p>	
<p><b>(v)</b></p>	<p>Let <math>W</math> be the number of patients who require surgery out of 15.  <math>W \sim \text{B}(15, 0.4)</math>  <math>E(W) = 15(0.4) = 6</math>  <math>\text{Var}(W) = 15(0.4)(1 - 0.4) = 3.6</math></p>	
	<p>Let <math>\bar{W}</math> be the mean number of patients who require surgery.  Since <math>n</math> is large, so by Central Limit Theorem,  <math>\bar{W} \sim \text{N}\left(6, \frac{3.6}{50}\right)</math> approximately.  <math>P(\bar{W} &gt; 5.5) = 0.96880</math>  <math>\approx 0.969</math> ( to 3 s.f )</p>	